

Discoloration of Teeth after Avulsion and Replantation: Results from a Multicenter Randomized Controlled Trial

Peter F. Day, PhD, FDS,* Monty S. Duggal, PhD, FDS,* Alec S. High, PhD, FRCPath,* Angus Robertson, BSc, FIMI,* Terry A. Gregg, FDS, FFD,[†] Paul F. Ashley, PhD, FDS,[‡] Richard R. Welbury, PhD, FDS,^{||} Ben O. Cole, MSc, FDS,[¶] and Stephen Westland, PhD, CCol FSDC[#]

Abstract

Introduction: There is evidence to suggest that Ledermix, placed as an intervisit root canal dressing, might improve periodontal healing after replantation of avulsed teeth. As a part of a multicenter randomized controlled trial, we aimed to compare the effect of 2 root canal medicaments, Ledermix and Ultracal XS, on the discoloration of replanted teeth. **Methods:** Discoloration was investigated by using 3 methods: patient satisfaction with the color of replanted teeth, clinical photographs taken at baseline and 12-month reviews, and estimation of color change by using CIELAB scores for baseline and 12-month photographs. **Results:** Twenty-two patients (27 teeth) were recruited. Ten patients (12 teeth) were randomized to the Ledermix group and 12 patients (15 teeth) to the Ultracal XS group. At 12 months, 8 patients were concerned with the discoloration of their teeth. Seven came from the Ledermix group and 1 from the Ultracal XS group. This difference was significant (Fisher exact test, $P = .009$). Standardized photographs were taken for the patients recruited at one center only (17 patients). There was significant discoloration of teeth from baseline with Ledermix, causing a darkening and gray-brown discoloration (mean change from baseline to 12 months, $L^* = -5.1$, $a^* = 0.3$, $b^* = -1.2$, and $\Delta E = 8.1$) and Ultracal XS, causing a yellowing and lightening of teeth ($L^* = 1.9$, $a^* = 0.3$, $b^* = 3.3$, and $\Delta E = 5.4$). There was a significant difference for the L^* and b^* variables (independent t test) between the 2 groups. **Conclusions:** Both root canal medicaments cause discoloration, with Ledermix proving less acceptable to patients. (*J Endod* 2011;37:1052–1057)

Key Words

Avulsion, discoloration, Ledermix, multi-center, non-setting calcium hydroxide, randomized controlled trial

Discoloration of teeth after replantation after avulsion is recognized. Ebeleseder et al (1) reported a prevalence of severe discoloration of 33% for immature teeth that had been given the opportunity to revascularize. Another study reported that 27 of 28 teeth showed adverse color changes ranging from grayish to bluish-black (2).

Although color change has been reported as a consequence of dentoalveolar trauma by a number of authors, the mechanisms are not fully understood (3, 4). It does appear that different colors represent different clinical entities, some of which represent pulpal damage with subsequent healing, whereas others indicate pulpal necrosis. Furthermore, discoloration can occur as a result of materials used for subsequent endodontic treatment (5, 6).

Ledermix (Henry Schein UK [Holdings Limited]) has demonstrated increased rates of periodontal healing in a number of animal studies (7–11) for avulsed and replanted teeth. Ledermix is a water-soluble paste containing 1% triamcinolone, a corticosteroid, and 3% demeclocycline, a tetracycline. Two *in vitro* studies have investigated the effect of discoloration with Ledermix in immature (12) and mature roots (13). These studies showed that Ledermix caused a dark, gray-brown discoloration of teeth that was exacerbated by sunlight and immature root development. When Ledermix was restricted to the root with no material allowed into the crown, the degree of discoloration was less severe.

After avulsion and replantation in closed apex teeth, current guidelines (14) advocate that pulp extirpation and disinfection are instigated at day 7–10. The reason for this is that the recommended intracanal medicament, non-setting calcium hydroxide, showed significant detrimental outcomes to periodontal healing in comparison to a more inert gutta-percha root filling if placed earlier (15, 16). There are a number of studies in the literature with respect to color change after application of calcium hydroxide (12, 13, 17). It has also been noted that a more significant yellow color change was associated with immature rather than mature teeth (12).

Color change in teeth is difficult to measure because of many different factors. These include the effect of adjacent tissues (eg, gingivae), the area of tooth being measured, surface imperfections and defects, lighting conditions, angulations of the tooth surface, and dehydration. Consequently, a number of methods have been established and tested to determine tooth color or color changes over time (18). The simplest method is evaluating the tooth next to a shade guide of different tooth colors to monitor changes. Other methods include photographing the teeth in standardized conditions by using an adjacent universal gray block or the use of

From the *Leeds Dental Institute, Leeds; [†]Belfast Children's Hospital, Belfast; [‡]UCL Eastman Dental Institute, London; ^{||}Glasgow Dental Hospital, Glasgow; [¶]Newcastle Dental Hospital, Newcastle Upon Tyne; and [#]School of Design, University of Leeds, Leeds, United Kingdom.

This study was partially funded by a Department of Health grant and Henry Schein, who purchased specific equipment (5 Periostests). Free materials for the trial were provided by Mediartis, TTS splints; Henry Schein, Ledermix; and Optident, Ultracal XS.

Address requests for reprints to Dr Peter F. Day, Department of Paediatric Dentistry, Leeds Dental Institute, University of Leeds, UK LS2 9JT. E-mail address: p.f.day@leeds.ac.uk

0099-2399/\$ - see front matter

Copyright © 2011 American Association of Endodontists.

doi:10.1016/j.joen.2011.04.014

expensive spectrophotometers or colorimeters. All these methods have their disadvantages.

However, apart from shade guides, these other systems have one distinct advantage in that they can transform the results into a numeric color scale, CIE (1976) $L^*a^*b^*$ (abbreviated to CIELAB). CIELAB represents an approximately uniform color space, or globe, with equal distances corresponding to near-identical perceived color differences (18). The advantage of using this system is that for each color, a numeric value is given for 3 coordinates. In addition, where there is a change in color with time or objects of 2 different colors, the change can be calculated for the 3 coordinates (ΔL^* , Δa^* , Δb^*), and the distance between 2 points on the color space can be calculated to give a ΔE (Fig. 1). Two clinical studies (19, 20) investigated the size of the color mismatch needed between teeth, ΔE 2.6 and ΔE 3.7, before this was identified by dentists. A difference of ΔE 5.5 and 6.8 was required for dentists to find the color mismatch sufficiently unacceptable to suggest further treatment to improve it.

The aim of this study was to investigate the discoloration of teeth after application of 2 root canal medicaments for replanted teeth by using a multicenter randomized controlled trial (RCT).

Methods

This article reports on one of the secondary outcome measures for a multicenter RCT designed to study outcomes of teeth, replanted after avulsion, in which the initial root canal medicament was either Ultracal XS (a commercially prepared non-setting calcium hydroxide paste with high viscosity; Ultradent Products Inc, South Jordan, UT) or Ledermix. The primary outcome measure was periodontal healing; this has been reported in a different article (21). Before the study could commence a number of research approvals were secured, including ethical approval (04/Q1205/145), Medicines and Healthcare products Regulatory Agency (MHRA) (CTA 16767/0203/001-0001), European Union Drug Regulating Authorities Clinical Trials (EudraCT) number (2004-005188-12), and registration of the trial with International Standard Randomized Controlled Trial Number (ISRCTN) (58467151). Informed consent was obtained from all patients recruited to the study. All avulsed teeth were replanted within 60 minutes. The trial medicaments were placed in the root canal within 10 days (baseline) and remained in situ for 60–90 days. Both groups were then subject to an additional 3 months of intracanal dressing with Ultracal XS before obturation with gutta-percha and Tubliseal (SybronEndo, Orange, CA) at 6 months. Patients were reviewed at 12 months. At this appointment the patients were asked whether they were happy with the color of the avulsed and replanted tooth or teeth.

All patients who had care provided at Leeds Dental Institute had photographic records taken at each clinical visit to monitor discoloration. These were taken in the photography department by trained medical photographers with a Fujifilm (Tokyo, Japan) Finepix S3 Pro camera with 105-mm Micro Nikon (Tokyo, Japan) f2.8 lens and sigma ring flash, EM-140DG. Although attempts were made to standardize the way all photographs were taken, there was some variability in each image, depending on the camera used, the lighting, the photographer, or the photographic view taken.

To facilitate comparison between photographs and then quantify any color change, dedicated software was developed by using MathWorks (Natick, MA). The software, which is described in more detail in Figure 2, calculated the red-green-blue (RGB) and L^* , a^* , b^* values for 2 areas of interest, after it had first corrected the color of the second photograph to that of the first. The program then displayed these values for the reference (baseline) and




Figure 1. The 3-dimensional color space CIE (1976) $L^*a^*b^*$ (18). For every color a point in this space can be identified by using the 3 color coordinates L^* , a^* , and b^* . A change in color (from point X to X_1) can be calculated numerically by the distance between the 2 points ΔE . This value will only give the size of the color change, not the direction. For this the individual ΔL^* , Δa^* , Δb^* would be needed.

corrected (12-month) photographs, together with the change in color, CIELAB ΔE .

The dedicated software (Fig. 2) was also used to estimate the difference in color between 2 different teeth in the same photograph. By fixing the ratio for color correction so that none occurred, the same photograph could be imported into the upper left, upper right, and lower right panels. This then allowed the avulsed tooth and contralateral nonavulsed tooth to be compared in the 12-month photograph.

The L^* , a^* , b^* values for the baseline and 12-month photographs (including contralateral nonavulsed teeth) together with the CIELAB ΔE were recorded onto an SPSS spreadsheet (Statistical Package for the Social Sciences, version 16.1; SPSS Inc, Chicago, IL). Continuous data were assessed for whether they were normally distributed by using a Shapiro-Wilks test, and this determined whether an independent t or Mann-Whitney U test was used. A summary statistic, change in $L^*a^*b^*$, was calculated for each avulsed tooth between the baseline and 12-month photographs. The statistical tests were applied to these data. For dichotomous patient satisfaction data, either Pearson χ^2 or Fisher exact test was used, dependent on the numbers of observations in each cell.

Results

Eight of the 22 patients recruited at all centers were concerned with the color of their avulsed teeth. Seven of these patients had been randomized to the Ledermix group ($n = 10$). One of these patients was concerned with the color of 1 tooth but not with 2 other replanted teeth. One patient ($n = 12$) in the Ultracal XS group expressed dissatisfaction with the color. This difference was significant when using a Fisher exact test, whether the patient with multiple teeth in the Ledermix group was included in both groups ($P = .009$) or excluded ($P = .02$) from the analysis.

For the patients who were seen at Leeds Dental Institute, samples of the baseline and final 12-month review photographs are shown in Figure 3 before color correction. There was




Figure 2. Dedicated software was used to correct the color of photographs taken at different visits to then calculate whether there was any change in color between the 2 time points or between the 2 treatment groups. The picture in the top left panel was the original photograph taken at the initial visit (baseline). The picture in the top right panel was a picture taken at a subsequent review visit (12-month). Before any change in color could be established between the two, the color of the 12-month photograph was corrected to that of the baseline photograph. This was carried out by identifying a color stable, chromatically neutral, reference point in both photographs, for example, an uninjured adjacent tooth. In this case part of tooth #11 was used. The program calculated the XYZ values (this is an intermediary space between RGB and CIE LAB) for this reference point between the 2 photographs and then used this ratio to correct the color in the second photograph back to that of the first. The color-corrected 12-month picture was then shown in the lower right panel.

a noticeable darkening and graying effect of the teeth in the Ledermix group and a yellowing effect in the Ultracal XS group. Furthermore, there was variation in color within each root canal medicament group.

The change in color between baseline and 12 months was quantified for each tooth by using the dedicated software. Twenty-two teeth (17 patients) seen in Leeds Dental Institute were evaluated. For 2 patients in the Ledermix group, the data set was incomplete. For 1 patient with 1 avulsed tooth, a baseline and day 60–90 photographs were all that were available, because this patient failed to attend further appointments. Therefore, the results from the day 60–90 photograph were grouped with 12-month data. For the second patient with 1 avulsed tooth, no baseline photograph was taken; therefore, the data from the day 60–90 were analyzed with baseline photograph data. Table 1 shows baseline, 12-month, and change in color for L*, a*, b* in the CIE LAB space and CIE LAB ΔE values. There was a significant difference for L* ($P = .003$; confidence interval, 2.6–11.3; independent t test) and b* ($P = .009$; confidence interval, 1.3–7.9; independent t test). The Ultracal XS baseline data for a* were not normally distributed as determined by the Shapiro-Wilks test for normality. Therefore, for these

data the median and interquartile range are reported. There was no significant difference for the change in color between the groups as determined by Mann-Whitney U test. At baseline, day 0–10, there was no significant difference between Ledermix and Ultracal XS groups with respect to L*a*b* (independent t test or Mann-Whitney U test, depending on normality of data).

To validate this change in color between the 2 groups, where a contralateral central incisor that had not been avulsed was available, the difference in color between the 2 teeth was calculated for the 12-month review photograph. For this, only 12 teeth (12 patients) were eligible, with 7 from the Ledermix group and 5 from the Ultracal XS group. One patient in the Ledermix group had only the day 60–90 photograph, because the patient was lost to follow-up. Therefore, this photograph was used for comparison. This patient and another who had avulsed canines had their uninjured contralateral canine used for comparison. The change in color between the 2 teeth is displayed in Table 2. There was a significant difference for L* ($P = .007$; Mann-Whitney U test) variable and CIE LAB ΔE value ($P = .045$; independent t test; confidence interval, 0.12–8.82). For the contralateral nonavulsed tooth there was no significant difference between the Ledermix and

ULTRACAL XS GROUP



Baseline v. 12 months
Tooth avulsed #9
Not concerned about colour



Baseline v. 12 months
Teeth avulsed #8, #9
Concerned about colour

LEDERMIX GROUP



Baseline v. 12 months
Tooth avulsed #9
Concerned about colour



Baseline v. 12 months
Tooth avulsed #8
Not concerned about colour

Figure 3. The baseline and 12-month review photographs for patients seen at Leeds Dental Institute. The avulsed and replanted teeth were treated as part of a multicenter RCT, with teeth randomized to Ultracal XS or Ledermix root canal medicaments. Each photograph details the avulsed and replanted teeth and whether the patient was concerned with the color. Two cases from each group have been selected. These examples show the range of discoloration seen.

Ultracal XS groups when using appropriate statistics (independent *t* test or Mann-Whitney *U* test, depending on normality of data).

Discussion

This study used 3 methods to assess discoloration: patient satisfaction with the color of replanted teeth, clinical photographs taken at baseline and 12-month reviews, and estimation of color change by using CIELAB scores for baseline and 12-month photographs.

It is important to investigate whether children were happy with the color of their avulsed and replanted teeth. In adult studies, dissatisfaction with the color of their teeth is a common complaint and reason to seek dental care (22). This was especially true for women and younger people (23). For children, the consequences of poor dental appearance

or untreated dental trauma led to negative social judgments being made by their peers (24, 25). Dental features have been shown to be one of the most common causes for teasing at school (26). Consequently, for children and adolescents involved in this multicenter RCT, the appearance of their teeth was important, and for some this might have influenced their own self-perception and sense of how others perceived them. Figure 3 shows examples of the difference in baseline and 12-month photographs for both medicaments. Patient dissatisfaction with the color of their replanted teeth was more common in the Ledermix group. This was a significant finding, despite the small number of patients recruited. What is interesting is that some children with minor discoloration were unhappy with the color of their teeth, when others with more marked discoloration were not. This phenomenon has been investigated with children who have enamel opacities. Different

TABLE 1. The Color of Avulsed Teeth at Baseline and 12 Months and the Color Change That Took Place during This Period

Color variables	Randomization	No. of teeth	Mean at baseline (SD)	Mean at 12 months (SD)	Mean change (SD), 12 months–baseline
L* ++	Ledermix	10	58.9 (8.0)	53.8 (10.3)	–5.1 (5.8)
	Ultracal XS	12	56.2 (6.9)	58.1 (6.8)	1.9 (3.8)
a* (+)	Ledermix	10	4.5 (3.6)	6.0 (6.4)	0.3 (5.5)
	Ultracal XS	12	3.6 (3.3)	4.7 (3.9)	0.3 (3.1)
b* ++	Ledermix	10	12.1 (5.1)	10.9 (6.8)	–1.2 (4.8)
	Ultracal XS	12	11.3 (3.0)	14.9 (2.8)	3.3 (2.4)
CIELAB ΔE	Ledermix	10			8.12 (4.9)
	Ultracal XS	12			5.4 (3.1)

All injured teeth were avulsed and replanted as part of multicenter RCT. For each variable (L*, a*, b* in the CIELAB space and CIELAB ΔE) the data were separated into the 2 treatment groups the teeth were randomized to. CIELAB ΔE is a summary value that used data from each of the L*, a*, b* variables and reports the distance in a 3-dimensional model of color between the baseline and 12 months. This value only quantifies change of color but not the direction. The variables (L*, a*, b*) report the individual component's color (white–black, green–red, and yellow–blue).

SD, standard deviation.

++Significant difference between the Ledermix and Ultracal XS groups for L* and b* by using an independent *t* test.

(+)Data are not normally distributed; therefore, the values for a* are given as a median and interquartile range.

children's perceptions of themselves and their friends varied according to the importance they placed on different domains such as physical attributes or personality (27).

Although there are many techniques to monitor color change, these frequently use expensive equipment and require a fixed jig for each patient to ensure the assessment is made from an identical point. Therefore, the method this study used allowed standardized photographs taken by professional medical photographers to be used. This minimized the time required for this part of the clinical visit, when other clinical and radiographic special tests were required in addition to providing the treatment. The methodology used to correct the photographs taken in medical photography allowed quantitative assessment of color change, in comparison to simply looking at the baseline and 12-month photographs. There were 2 processes when using the dedicated software, which could account for error. The first was the identification of a color stable point on an uninjured tooth that was used to standardize the color in the 12-month photograph to that of the baseline photograph. The second was the failure to identify the same area of interest in both photographs. The size of each error was estimated and could account for up to a CIELAB ΔE value of 1. Therefore, the errors within the dedicated software were up to CIELAB ΔE value of 2.

With respect to the significant outcomes, a cautionary comment should be made about the statistics used for the longitudinal assessment of color change. For 3 patients (7 teeth), each tooth is not an independent variable, because they are subject to the same host responses;

therefore, the estimation of effect would be more accurately calculated by using multilevel modeling statistics. However, this is a time-consuming process, and the size of the effect, which is clearly visible in Figure 3, might be able to compensate for the simple statistics used. These statistical issues do not extend to the comparison with the nonavulsed contralateral tooth because these 3 patients had no such tooth to make the comparison and were therefore excluded from the analysis.

For the Ledermix group there was a mean change in color from baseline to 12 months of L* = –5.1 (the tooth got darker), a* = 0.3 (tooth got redder), and b* = –1.2 (the tooth got bluer). For all these changes the standard deviation was larger than the mean change, showing a wide range in color changes seen. Values from *in vitro* work for both mature and immature teeth confirmed this direction and the approximate magnitude of color change including the insignificant finding for a* variable (12, 13). The methodology used, multicenter RCT, was the most robust way of proving an association because potential confounders, either known or unknown, had a random chance of being allocated to either of the study groups. Comparison of baseline photographs taken within 10 days of the injury of the nonavulsed contralateral tooth showed no significant difference between the groups, which demonstrates further the benefit of randomization. Which constituent within Ledermix is the cause of the discoloration is still unproven, because no results with regard to color were provided by the one animal study that

TABLE 2. The Change in Color Variables between an Avulsed Tooth Treated as Part of a Multicenter RCT and the Contralateral Nonavulsed Tooth

Color variables for CIELAB space	Randomization	No. of teeth	Avulsed tooth, mean (SD)	Nonavulsed contralateral tooth, mean (SD)	Difference between avulsed tooth and contralateral nonavulsed tooth (SD)
L* ++ (+)	Ledermix	7	62.4 (20.0)	54.4 (16.8)	–5.3 (6.8)
	Ultracal XS	5	67.9 (10.6)	67.1 (13.1)	0.3 (3.0)
a*	Ledermix	7	6.4 (3.0)	5.4 (3.8)	1.0 (1.8)
	Ultracal XS	5	3.9 (1.4)	4.1 (1.6)	–0.2 (0.3)
b*	Ledermix	7	13.2 (7.3)	12.9 (5.4)	0.2 (3.8)
	Ultracal XS	5	13.9 (3.8)	11.5 (3.9)	2.4 (0.6)
CIELAB ΔE ++	Ledermix	7			7.5 (4.2)
	Ultracal XS	5			3.0 (1.3)

For central incisors (n = 10) this was compared with the nonavulsed central incisor; for canines (n = 2) this was compared with the contralateral canine. For each color variable (L*, a*, b* in the CIELAB space and CIELAB ΔE) the data were separated into the treatment the teeth received. CIELAB ΔE is a summary value that used data from each of the L*, a*, b* variables and reports the distance between the avulsed and nonavulsed tooth in a 3-dimensional model of color. This variable only quantifies the change of color but not the direction. The variables (L*, a*, b*) report the individual component's color (white–black, green–red, and yellow–blue).

RCT, randomized controlled trial; SD, standard deviation.

++Significant difference between the Ledermix and Ultracal XS groups for L* by using a Mann-Whitney *U* test and CIELAB ΔE by using an independent *t* test.

(+)Data are not normally distributed; therefore, the values for L* are given as a median and interquartile range.

Acknowledgments

The authors would like to thank all staff at each center who helped with the running of this trial.

The authors deny any conflicts of interest related to this study.

References

- Ebeleseder KA, Friehs S, Ruda C, Pertl C, Glockner K, Hulla H. A study of replanted permanent maxillary incisors after replantation related to age and sex. *Dent Traumatol* 1998;14:274–8.
- Ravn JJ, Helbo M. Replantation of akcidentelt eksarticulerede tinder. *Tandloegbladet* 1966;90:805–15.
- Andreasen JO, Andreasen FM, Andersson L. Textbook and colour atlas of traumatic injuries to the teeth. 4th ed. Copenhagen: Blackwell Munksgaard; 2007.
- Holan G, Fuks AB. The diagnostic value of coronal dark-gray discoloration in primary teeth following traumatic injuries. *Pediatr Dent* 1996;18:224–7.
- Kupietzky A, Holan G. Treatment of crown fractures with pulp exposure in primary incisors. *Pediatr Dent* 2003;25:241–7.
- Plotino G, Buono L, Grande NM, Pameijer CH, Somma F. Nonvital tooth bleaching: a review of the literature and clinical procedures. *J Endod* 2009;34:394–407.
- Bryson EC, Levin L, Banchs F, Abbott PV, Trope M. Effect of immediate intracanal placement of Ledermix paste on healing of replanted dog teeth after extended dry times. *Dent Traumatol* 2002;18:316–21.
- Chen H, Teixeira FB, Ritter AL, Levin L, Trope M. The effects of intracanal anti-inflammatory medicaments on external root resorption of replanted dog teeth after extended extra oral dry time. *Dent Traumatol* 2008;24:74–8.
- Pierce A, Lindskog S. The effect of an antibiotic corticosteroid combination on inflammatory root resorption. *in vivo*. *Oral Surg Oral Med* 1987;64:216–20.
- Thong YL, Messer HH, Siar CH, Saw LH. Periodontal response to two intracanal medicaments in replanted monkey incisors. *Dent Traumatol* 2001;17:254–9.
- Wong KS, Sae-Lim V. The effect of intracanal Ledermix on root resorption of delayed-replanted monkey teeth. *Dent Traumatol* 2002;18:309–15.
- Kim ST, Abbott PV, McGinley P. The effect of Ledermix paste on discoloration of immature teeth. *Int Endod J* 2000;33:233–7.
- Kim ST, Abbott PV, McGinley P. The effect of Ledermix paste on discoloration of mature teeth. *Int Endod J* 2000;33:227–32.
- Flores MT, Andersson L, Andreasen JO, et al. Guidelines for the evaluation and management of traumatic dental injuries: II—avulsion of permanent teeth. *Dent Traumatol* 2007;23:130–6.
- Andreasen JO, Kristerson L. The effect of extra-alveolar root fillings with calcium hydroxide on periodontal healing after replantation of permanent incisors in monkeys. *J Endod* 1981;7:349–54.
- Lengheden A, Blomlof L, Lindskog S. Effect of immediate calcium hydroxide treatment and permanent root-filling on periodontal healing in contaminated replanted teeth. *Scand J Dent Res* 1990;99:139–46.
- Tinaz AC, Kivanc BH, Gorgul G. Staining potential of calcium hydroxide and monochlorophenol following removal of AH26 root canal sealer. *J Contemp Dent Pract* 2008;9:56–63.
- Joiner A. Tooth colour: a review of the literature. *J Dent* 2004;32:3–12.
- Douglas RD, Steinhauer TJ, Wee AG. Intraoral determination of the tolerance of dentists for perceptibility and acceptability of shade mismatch. *J Prosthet Dent* 2007;97:200–8.
- Johnston WM, Kao EC. Assessment of appearance match by visual observation and clinical colorimetry. *J Dent Res* 1989;68:819–22.
- Day PF, Gregg TA, Ashley P, et al. Periodontal healing following avulsion and replantation of teeth: a multi-center randomised controlled trial. *Dent Traumatol*. In press.
- Tin-Oo MM, Saddki N, Hassan N. Factors influencing patient satisfaction with dental appearance and treatments they desire to improve aesthetics. *BMC Oral Health* 2011; 11:6. Available at: <http://www.biomedcentral.com/1472-6831/11/6>. Accessed May 26, 2011.
- Vallittu PK, Vallittu ASJ, Lassila VP. Dental aesthetics: a survey of attitudes in different groups of patients. *J Dent* 1996;24:335–8.
- Rodd HD, Barker CS, Baker SD, Marshman Z, Robinson PG. Social judgements made by children in relation to visible incisor trauma. *Dent Traumatol* 2010;26:2–8.
- Bos A, Hoogstraten J, Prahl-Andersen B. Dutch primary school children's attitudes towards their dental appearance. *Pediatr Dent* 2008;30:439–42.
- Shaw WC, Meek SC, Jones DS. Nicknames, teasing, harassment and the salience of dental features among school children. *Br J Orthod* 1980;7:75–80.
- Marshman Z, Gibson B, Robinson PG. The impact of developmental defects of enamel on young people in the UK. *Community Dent Oral* 2008;37:45–57.
- Andreasen FM. Transient apical breakdown and its relation to color and sensibility changes. *Dent Traumatol* 1986;2:9–19.
- Joiner A, Hopkinson I, Deng Y, Westland S. A review of tooth colour and whiteness. *J Dent* 2008;36s:2–7.

Conclusions

Ledermix caused a significantly greater level of patient dissatisfaction with the color of avulsed and replanted teeth.

During a 12-month period Ledermix caused a significant gray-brown darkening of avulsed and replanted teeth.

During a 12-month period Ultracal XS caused a significant yellowing of the avulsed and replanted teeth.

In comparison to nonavulsed contralateral tooth, Ledermix caused a significant darkening of the tooth, whereas Ultracal XS caused yellowing, but this was not significant.